

Influence of different plant densities on crop yield of six safflower genotypes under Egyptian newly reclaimed conditions

Mohamed Hamza

Agronomy Department, Faculty of Agriculture, Cairo University, Cairo, Egypt

Corresponding author email: mhamza4000@gmail.com

ABSTRACT: Safflower (*Carthamus tinctorius* L.) is the crop of small holder farmers and its cultivation is available only in newly reclaimed soil. Density of cultivation in these lands should be tested in a random sample of genotypes. Thus, six safflower genotypes were evaluated in the Desert Experiment Station of the Faculty of Agriculture, Cairo University in Wadi El-Natroon, El-Beheira Governorate, Egypt in 2011/2012 and 2012/2013 seasons. Six safflower genotypes were tested under three plant densities (80000, 160000 and 240000 plant ha⁻¹) in a split-plot design. The main plots were devoted to the plant densities and sub-plots to the six genotypes were to facilitate the genetic dissection of salt-stress adaptive traits. Plant density of 240000 plants ha⁻¹ recorded the highest values of plant height, seed and oil yield kg ha⁻¹. Meanwhile, the highest values of number of branches and capitula, petal yield, seed yield plant⁻¹ were recorded at 80000 plant ha⁻¹. Line-1697 and Demo-137 surpassed the other genotypes in seed and oil yield kg ha⁻¹. Plant densities vs. genotypes interaction was significant for number of capitula, petal and seed yields plant⁻¹, seed index, seed oil %, seed and oil yield kg ha⁻¹. The highest seed yield (2890 kg) was realized from Line-1697 with 240000 plant ha⁻¹. Meanwhile, the highest oil yield (927.2 kg) was achieved with 240000 plant ha⁻¹ by Demo-137. Phenotypic and genotypic variances among lines were highly significant.

Key words: *Carthamus tinctorius*, Safflower, Genotypes, Density, Phenotypic variance, Genotypic variance, Broad sense heritability.

INTRODUCTION

Safflower is more drought resistance than other oilseeds and can produce good yield in dry region, while its salt tolerance is a valuable asset as the area affected by some degree of salinity steadily increases world-wide (Weiss, 2000). Safflower became a major oilseed in many countries in the world like USA especially after 2nd world war when breeders produced high yielding, high oleic oil cultivars with greater resistance to biotic and abiotic stresses. But it suffered severally from competition main winter crops in Egypt, i.e., berseem and wheat. Still the local edible oil production, in Egypt, is not sufficient. Safflower may play an important role in increasing edible oil production in new reclaimed soils in Egypt (Abu-Hagaza et al., 2009). The main obstacles for planting safflower are the limited genotypic materials and their narrow genetic background with respect to adaptation to these conditions in Egypt. Weiss (2000) mentioned that determination of the optimum plant population for a particular area under specific cropping condition is essential to optimize yield.

Plant densities are needed to be accurately established. The present stand (number of plants) of safflower may be insufficient to produce high yield via land races and the exotics, since a considerable number of plants may be lost during the growing season. Therefore, closer plant spacing could be more suitable to compensate for this loss and insure higher yield.

Many researchers reported that plant densities had significant effect on safflower yield and its components. Yau (2009) found that plant height decreased by increasing plant density. Amoughin et al (2012a), Sharifmghaddasi and Omidi (2009) found that increasing plant density increased plant height. By contrast, number of branches m⁻² decreased. Number of capitula plant⁻¹ or m⁻² decreased by increasing the plant density (Amoughin et al., 2012 a, Elfadl et al., 2009, Emami et al., 2011, Vaghar et al., 2014, Sharifmghaddasi and Omidi 2009). Besides, Shahri et al (2013) reported that increasing plant density increased number of capitula per unit area but decreased seed index. Conversely, Emami et al (2011), Sharifmghaddasi and Omidi (2009) mentioned that 1000-seed weight was increased by increasing plant

density. Ghanem and Ash-Shormillesy (2007), in Egypt, reported that seed yield plant⁻¹ decreased by increasing plant densities. They reported considerable variations in seed index (1000-seed weight) due to plant densities.

Amoughin et al (2012 b), Shahri et al (2013) and Yau (2009) reported that increasing plant density decreased oil %. But, Emami et al (2011) found that increasing plant density increased oil %. In the context, Amoughin et al (2012 b), Elfadl et al (2009), Emami et al (2011), Vaghar et al (2014), Sharifmghaddasi and Omid (2009) and Yau (2009) found that increasing plant density increased seed and oil yield ha⁻¹. On the other hand, Sharifi et al (2012) mentioned that increasing plant density decreased seed and oil yield ha⁻¹.

Yield and yield components of safflower were significantly affected by the genotypes (Abu-Hagaza 1990, Abu-Hagaza et al., 2009, Camas et al., 2007, Elfadl et al., 2009, Eslam 2004, Mokhtassi-Bidgoli et al., 2007, Sharifi et al., 2012, and Muñoz-Valenzuela et al., 2007 and Vaghar et al., 2014). In contrast, Samanci and Ozkaynak (2003) found no significant differences among safflower genotypes in seed yield. Meanwhile, they recorded significant differences among genotypes in oil content. Sharifi et al (2012) reported that the interaction between genotypes and plant densities was significant.

Camas and Esendal (2006) found that the heritability for plant height, number of branches, seed yield, 1000-seed weight and oil content were estimates as 93%, 45%, 35%, 81% and 59%, respectively. The plant height was the least affected trait over environments and followed by 1000-seed weight. On the other hand, seed yield, number of branch, and oil content were the most affected traits versus environmental conditions.

The objectives of the present research were to examine the performance of a random sample of exotic and landraces of safflower genotypes under three plant densities to determine the optimum plant density for each genotype and to estimate some genetic parameters of these genotypes under the stress condition of the newly reclaimed soil of Wadi El-Natroon and the different densities used herein.

MATERIALS AND METHODS

Two field experiments were carried out at the Agricultural Experiments Desert Station, Faculty of Agriculture, Cairo University in Wadi El-Natroon, El-Beheira Governorate, during the two winter seasons of 2011/2012 and 2012/2013 under drip irrigation system. The soil of the experimental site was sandy (93% sand) and saline (7.5 Ec; ds/m) with a pH of 7.8. Also, the irrigation water is saline (4.2 Ec; ds/m) according to the analysis of soil and water.

A split-plot design, in randomized complete blocks arrangements, with three replications was used. The main plots were devoted to three plant densities, viz., 84, 168 and 252 thousand plants ha⁻¹. The sub-plots were allotted to a random sample of six safflower genotypes that represents land races and exotics beside the local safflower cultivar. Three of these genotypes, were local entries including Giza-1, the sole cultivar grown in Egypt, and 2 land races collected from 2 Governorates as farmer's seed lots, i.e. Bani-Suef (Somosta center) and Aswan (Daraw center) representing middle and upper Egypt, respectively. The remaining 3 exotics viz. Demo-137 cv. from USA, Line-1697 from Cyprus and Line-168 from Turkey. All seeds were kindly devoted by Oil Seed Crops Research Program, Field Crops Research Institute, Agricultural Research Center (ARC), Ministry of Agriculture, Egypt.

Each sub-plot consisted of 5 rows of 4 m length and 0.60 m width. The experimental plot area was 12 m². Seeds were sown in hills 20, 10 and 7 cm apart on 15 October in both seasons, thereafter were thinned to one plant hill⁻¹ to give the three plant densities (80000, 160000 and 240000 plant ha⁻¹).

Single super-phosphate fertilizer (15.5% P₂O₅) at the rate of 72 kg P₂O₅ ha⁻¹ was applied uniformly before planting. Nitrogen was added at level of 180 kg N ha⁻¹, in the form of ammonium nitrate (33.5% N). The first dose was added at 21 days from planting, and then the rest doses were applied at a 7-day interval. Potassium sulphate (50% K₂O) at the rate of 120 kg K₂O ha⁻¹ was added through five equal doses at a 7-day interval. Other cultural practices were carried out in the proper time according to the package-deal of ARC.

At harvest, ten guarded plants were randomly sampled from the two inner rows of each sub-plot to determine plant height (cm), number of branches plant⁻¹, number of capitula plant⁻¹, petals yield plant⁻¹ (g), seed yield plant⁻¹ (g), seed index (100-seed weight g). Seed oil percentage was determined according to AOAC (2000). Seed yield (kg ha⁻¹) was weighed from the whole area of each sub-plot and adjusted to yield per hectare. Oil yield (kg ha⁻¹) was calculated by multiplying seed-oil percentage by seed yield ha⁻¹. All obtained data were statistically analyzed and means were compared by LSD test according to procedures outlined by Gomez and Gomez (1984) using MSTAT-C computer program (Freed et al., 1989). Test for homogeneity of variance was used to compare between variances over two years before deciding the validity of combined analysis.

The expected mean squares (EMS) shown in table (1) were used to estimate the genetic (σ_g^2) and genetic by year interaction (σ_{gy}^2) variances as follows according to Hallauer and Miranda (1988): $\sigma_g^2 = (M_3 - M_2)/ry$ and $\sigma_{gy}^2 = (M_2 - M_1)/r$, where r = number of replications and y = number of years. The phenotypic variance (σ_{ph}^2) was estimated as follows: $\sigma_{ph}^2 = \sigma_g^2 + (\sigma_{gy}^2 / r) + (\sigma_e^2 / ry)$. Heritability in the broad sense (h_b^2) was

estimated using the following formula: $h^2_b = (\sigma^2_g / \sigma^2_{ph}) \times 100$. Genetic coefficient of variability (G.C.V %) was

estimated as follows: $G.C.V = \sqrt{\frac{\sigma^2}{\bar{x}}} * 100$

Table 1. Expected mean squares (E.M.S) of combined analysis of variance across two years.

S.O.V.	df		M.S.	E.M.S.
Years (Y)	y-1	= 1		
Years/repes	Y (r-1)	= 4		
Genotypes (G)	g-1	= 5	M ₃	$\sigma^2_e + r \sigma^2_{gy} + ry \sigma^2_g$
G × Y	(g-1) (y-1)	= 5	M ₂	$\sigma^2_e + r \sigma^2_{gy}$
Pooled error	Y (r-1) (g-1)	= 52	M ₁	σ^2_e

RESULTS AND DISCUSSION

Analysis of variance

Combined analysis of variance (Table 2) showed that highly significant differences were existed among genotypes for all studied traits. Mean squares due to plant densities were also highly significant for all studied characters. Mean squares of years were insignificant for all traits except, number of branches plant⁻¹ and seed and oil yield ha⁻¹. They revealed highly significant differences in the two seasons. All mean squares due to genotypes × plant densities were highly significant for number of capitula plant⁻¹, seed index, seed yield plant⁻¹, seed and oil yields ha⁻¹. Meanwhile, difference was significant with petal yield plant⁻¹. Significant differences shown among plant densities × years for seed index, seed and oil yields ha⁻¹ only.

Genotypes × year's mean squares were insignificant for all traits except plant height, seed and oil yields ha⁻¹. Finally, mean squares for the second order interaction (genotypes × plant densities × years) were highly significant for petal yield plant⁻¹, seed and oil yields ha⁻¹ and significant for seed index. Confirming previous results (Amoughin et al., 2012 (a,b), Elfadl et al., 2009, Emami et al., 2011, Shahri et al., 2013, Sharifmghaddasi and Omidi 2009).

Effect of plant densities

Data in Table (3) revealed that plant height, number of branches plant⁻¹, number of capitula plant⁻¹, petal yield plant⁻¹, seed index, seed yield plant⁻¹, seed oil %, seed and oil yield ha⁻¹ were significantly affected by plant densities. Table (3) showed that plant density increased plant height up to 240000 plant ha⁻¹ due to reducing the light absorption inside the plant canopy and creating competition among plants. Similar trends were obtained by Amoughin et al (2012a), Sharifmghaddasi and Omidi (2009). Also, Yau (2009) found that plant height decreased by decreasing plant densities. Also, Table (3) showed that number of branches plant⁻¹ was decreased by increasing plant density up to 240000 plant ha⁻¹. Decreasing of plant density may be the main cause to increase light intensity around plants and encouraged branching. This result is in harmony with that of Amoughin et al (2012a) and of Sharifmghaddasi and Omidi (2009). Increasing number of capitula plant⁻¹ due to decreasing plant density was observed in Table (3), number of branches plant⁻¹ may be the main cause of this increase. Confirming previous results (Amoughin et al., 2012 (a), Elfadl et al., 2009, Emami et al., 2011, Vaghar et al., 2014, Sharifmghaddasi and Omidi 2009). In contrast, Shahri et al. (2013) reported that increasing plant density increased number of capitula. Results in the same table revealed that seed yield plant⁻¹ was increased by decreasing density levels up to 80000 plant ha⁻¹. Ghanem and Ash-Shormillesy (2007) reported similar result. Such increase may be attributed to the increase in number of branches and capitula plant⁻¹.

Combined data in Table (3), showed that decreasing density levels markedly increased seed index. Similar results were obtained by Shahri et al (2013). This increase may be explained by higher dry matter accumulation partitioned to seeds. Lower plant density increased also seed oil %. Same trend was observed by Amoughin et al (2012b), Shahri et al (2013), Yau (2009). However, Emami et al (2011) found that increasing plant densities increased seed oil %.

Table 2. Mean squares of combined analysis of variance for all studied traits of six safflower genotypes evaluated under three plant densities in 2011-12 and 2012-13 seasons

Source of variation	df	Mean of squares								
		Plant height (cm)	No. of branches plant ⁻¹	No. of capitula plant ⁻¹	Petal yield plant ⁻¹ (g)	Seed index (g)	Seed yield plant ⁻¹ (g)	Seed yield ha ⁻¹ (kg)	Seed oil (%)	Oil yield ha ⁻¹ (kg)
Years (Y)	1	22.69 ^{ns}	16.96 ^{**}	111.02 ^{ns}	0.08 ^{ns}	0.003 ^{ns}	81.23 ^{ns}	389859.23 ^{**}	0.99 ^{ns}	32968.32 ^{**}
R(Y)	4	128.10 [*]	1.57 ^{ns}	20.15 ^{ns}	0.43 ^{ns}	0.14 ^{ns}	16.83 ^{ns}	10393.23 ^{ns}	1.97 ^{ns}	601.73 ^{ns}
Plant densities (D)	2	856.04 ^{**}	151.03 ^{**}	1055.09 ^{**}	11.24 ^{**}	8.75 ^{**}	1017.46 ^{**}	721727.02 ^{**}	48.90 ^{**}	476813.07 ^{**}
YD	2	37.40 ^{ns}	1346 ^{ns}	25.52 ^{ns}	0.004 ^{ns}	0.53	9.80 ^{ns}	359353.57 ^{**}	0.02 ^{ns}	34552.52 ^{**}
Error (a)	8	22.48	0.57	53.60	0.11	0.08	21.75	5163.09	1.31	910.39
Genotypes (G)	5	1422.76 ^{**}	3.14 ^{**}	227.78 ^{**}	1.84 ^{**}	1.98 ^{**}	46.56 ^{**}	865111.65 ^{**}	94.40 ^{**}	257547.95 ^{**}
YG	5	108.09 [*]	0.24 ^{ns}	5.30 ^{ns}	0.31 [*]	0.18 ^{ns}	2.62 ^{ns}	153202.09 ^{**}	0.35 ^{ns}	16924.62 ^{**}
DG	10	61.37 ^{ns}	0.61 ^{ns}	58.36 ^{**}	0.27 [*]	0.54 ^{**}	35.32 ^{**}	230444.81 ^{**}	0.85 ^{ns}	25355.97 ^{**}
YDG	10	47.38 ^{ns}	0.68 ^{ns}	14.68 ^{ns}	0.32 ^{**}	0.31 [*]	9.74 ^{ns}	120945.14 ^{**}	0.1 ^{ns}	12651.52 ^{**}
Error (b)	60	39.21	0.80	17.53	0.11	0.14	5.96	22551.00	1.72	3609.68
C.V (%)		5.01	9.56	14.75	23.53	5.91	14.64	6.70	4.17	8.51

ns, * and ** indicate non-significance and significance at 5 and 1% probability levels, respectively.

Table 3. Effect of plant density and genotypes on safflower yields and its components (combined data of 2011-12 and 2012-13 seasons)

Trait	Plant height (cm)	No. of branches plant ⁻¹	No. of capitula plant ⁻¹	Petal yield plant ⁻¹ (g)	Seed index (g)	Seed yield plant ⁻¹ (g)	Seed yield ha ⁻¹ (kg)	Seed oil (%)	Oil yield ha ⁻¹ (kg)
Factor									
Plant density (D)									
80000	119.6	11.6	34.2	2.0	6.8	22.8	1749	32.6	576.5
160000	123.1	8.9	27.6	1.2	6.3	14.0	2351	31.5	743.0
240000	129.3	7.6	23.4	0.9	5.8	13.2	2624	30.3	797.5
L.S.D _{0.05}	2.6	0.4	4.0	0.2	0.5	2.5	39.0	0.6	16.4
Genotype (G)									
Giza-1	126.3	9.3	26.4	1.4	6.3	15.5	1982	29.6	579.5
Bani-Suef	135.8	9.1	26.1	1.2	6.0	15.9	2249	30.6	687.8
Aswan	130.0	8.9	25.0	0.9	5.9	15.2	1978	28.5	560.3
Demo-137	124.5	9.6	29.4	1.5	6.4	17.8	2386	33.6	798.5
Line -168	114.6	10.0	34.8	1.9	6.8	19.4	2510	34.3	859.7
Line -1697	112.7	9.5	28.7	1.4	6.4	16.3	2345	32.1	748.1
L.S.D _{0.05}	4.2	0.6	2.8	0.2	0.2	1.6	100.1	0.9	40.1

Table (3) showed that plant density increased seed yield ha⁻¹, such increase in seed yield may be attributed to the considerable increase in plant density. Amoughin et al (2012b), Elfadl et al (2009), Emami et al (2011), Vaghar et al (2014), Sharifmghaddasi and Omid (2009) and Yau (2009) stated the same results, meanwhile, Shahri et al (2013) mentioned opposite trend. With respect to oil yield, it was significantly increased by plant density (Table 3). This increase may be due to the increase in seed oil % and seed yield ha⁻¹. These results are in general agreement with those obtained by Amoughin et al (2012b), Elfadl et al (2009), Emami et al (2011), Vaghar et al (2014), Sharifmghaddasi and Omid (2009) and Yau (2009). On the contrary, Shahri et al (2013) cleared that oil yield was depressed significantly by increasing plant density.

Effect of genotypes

Combined data in Table (3) cleared that there were significant differences among all genotypes in plant height, number of branches plant⁻¹, number of capitula plant⁻¹, petal yield plant⁻¹, seed index, seed yield plant⁻¹, seed oil %, seed and oil yield ha⁻¹. Line-1697, recorded the shortest plant height (112.7 cm) when compared to the rest ones. Significant differences were recorded among genotypes concerning number of branches and capitula plant⁻¹. The exotic genotypes Line-168 followed by Demo-137 produced the highest number of branches and capitula plant⁻¹.

Data in Table (3) also revealed that both the exotic genotypes, Turkey-168 and Demo-137 had higher values of petals and seed yield plant⁻¹. Generally, this superiority may be due to the increase in number of branches and capitula plant⁻¹ and seed index. Concerning seed oil % Line-1697 and Demo-137 surpassed the rest of genotypes (Table 3). Samanci and Ozkaynak (2003) reported significant differences in oil content due to different genotypes.

In general, two promising exotics genotypes, namely, Line-168 and Demo-137 outyielded the other genotypes in seed yield ha⁻¹ (Table 3). This superiority may be a result of the increase in number of branches and capitula plant⁻¹, seed yield plant⁻¹ and seed index. These results are in harmony with those obtained by Camas et al (2007), Mokhtassi-Bidgoli et al (2007), Osman and Ali (2006), and Muñoz-Valenzuela et al (2007) who reported that there were significant differences among genotypes studied. On the other hand, Samanci and Ozkaynak (2003) found no significant difference among safflower genotypes used with respect to seed yield. Also, Osman and Ali (2006) reported that some introduced lines surpassed the commercial cultivar Giza-1 under modern system of irrigation in calcareous soils of Egypt.

Effect of the interaction

No significant interaction between plant density and genotypes used was detected for plant height, number of branches plant⁻¹ and seed oil %. Meanwhile, number of capitula plant⁻¹, petal and seed yield plant⁻¹, seed index and seed and oil yield ha⁻¹ significantly affected by plant populations.

Results in Table (4) showed the effect of the interaction between plant density and safflower genotypes on seed and oil yield ha⁻¹. Results revealed that the highest seed and oil yield was obtained from the interaction of Line-1697 × 240000 plant ha⁻¹ (2890, 900.3 kg, respectively) followed by Demo-137 × 240000 plant ha⁻¹ (2796, 927.2 kg, respectively) without significant differences. In the context, Sharifi et al (2012) reported that genotypes vs. plant densities interaction were significant.

Table 4. Effect of the interaction of plant density and genotype on seed and oil yields of safflower (combined data of 2011-12 and 2012-13 seasons).

Genotype (G)	Seed yield (kg ha ⁻¹)			Oil yield (kg ha ⁻¹)		
	Plant density (D) (10 ³ plant ha ⁻¹)					
	80	160	240	80	160	240
Giza-1	1416	2083	2446	444.1	613.7	680.5
Bani-Suef	1858	2460	2430	587.5	758.0	718.1
Aswan	1376	2098	2461	406.3	598.2	676.4
Demo-137	2346	2386	2796	840.3	811.7	927.2
Line -168	1919	2517	2723	661.0	852.4	882.2
Line -1697	1580	2565	2890	520.1	824.0	900.3
L.S.D _{0.05} (D×G)	173.4			69.38		

Genetic parameters

Changes in the magnitude of genotypic (σ_g^2) and phenotypic (σ_{ph}^2) variances, broad-sense heritability (h_b^2) and genetic coefficient of variability (G.C.V) of studied traits for the six genotypes under three plant densities across two years are presented in Table (5). In general, changes in magnitude of σ_g^2 and σ_{ph}^2 varied to plant densities environment for all traits.

Table 5. Genotypic (σ^2_g) and phenotypic (σ^2_{ph}) variances, heritability in the broad sense (h^2_b), grand mean and (G.V.C. %) for all studied traits under lower, normal and higher plant densities (data are combined across two seasons).

Trait variances	Plant height (cm)	No. of branches plant ⁻¹	No. of capitula plant ⁻¹	Petal yield plant ⁻¹ (g)	Seed index (g)	Seed yield plant ⁻¹ (g)	Seed yield ha ⁻¹ (kg)	Seed oil (%)	Oil yield ha ⁻¹ (kg)
Lower density			80000 plant ha ⁻¹						
σ^2_g	37.14	0.45	9.02	0.06	0.17	10.86	105728.73	5.16	21322.29
σ^2_{ph}	84.44	0.59	13.90	0.09	0.27	12.90	160939.25	5.56	28530.97
h^2_b (%)	43.99	75.60	64.93	63.65	63.86	84.22	65.69	92.80	74.73
Mean	119.63	11.62	34.15	6.79	2.01	22.80	1749.28	32.58	576.54
G.C.V (%)	5.10	5.78	8.78	3.45	20.81	14.46	18.59	6.97	25.33
Medium density			160000 plant ha ⁻¹						
σ^2_g	90.29	0.03	34.42	0.03	0	1.45	10458.75	5.06	9435.39
σ^2_{ph}	100.64	0.22	36.05	0.16	0.13	2.06	75388.37	5.23	14541.13
h^2_b (%)	89.72	15.22	95.48	16.42	0	70.26	13.87	96.85	64.89
Mean	123.08	8.88	27.56	6.30	1.22	14.04	2351.39	31.47	742.98
G.C.V (%)	7.72	2.06	21.26	2.54	0.00	8.59	4.35	7.14	13.07
Higher density			240000 plant ha ⁻¹						
σ^2_g	96.39	0	8.19	0.30	0.09	3.54	41045.84	5.71	13581.02
σ^2_{ph}	102.14	0.10	13.53	0.33	0.13	5.28	45390.30	6.00	14152.53
h^2_b (%)	94.37	0	60.52	90.70	69.23	67.05	90.43	95.08	93.96
Mean	129.26	7.61	23.43	5.81	0.93	13.20	2624.40	30.26	797.46
G.C.V (%)	7.59	0.00	12.23	9.43	34.13	14.26	7.72	7.88	14.61

Genotypic variance for plant height increased by increasing plant density. Thus, the differences among genotypes were obvious under the highest plant densities. Besides, heritability in the previous trait was medium and increased to high in the medium density and became very high in the highest density used. Thus, heritability is not a fixed number but apt to increase or decrease for different treatments. By contrast, No. of branches per plant showed limited phenotypic variation under the lowest density and no genotypic variation under 240000 plant ha⁻¹. In all cases for all traits and across all plant densities, phenotypic variance was higher than genotypic variance without any exception. High h² was high for seed yield per plant (84.2 %) in the lowest density and decreased with increasing density. Surprisingly, heritability for seed yield ha⁻¹ was difficult to explain. It was medium (65.69%) in the lowest density used with almost no difference due to genotype in the middle density and became very high (90.43%) under the highest density used. This contradictory could not be attributed to the behavior of genotypes used but to the soil of Wadi El-Natroon. Low to zero coefficient of variability was reported by number of branches per plant in the three densities used. The highest GCV % was observed for oil yield per hectare in the lowest density (25.33) followed by seed index (20.81) and seed yield ha⁻¹ (18.59). On the other hand, very high genetic coefficient was obtained for seed index (34.13). This contradictory in results is a result of the nature of the newly reclaimed soil. Thus, more experiments should be done at different location and in different seasons to have accurate estimation for different genetic parameters in newly reclaimed soils and especially at Wadi El-Natroon. Similar trends were observed by Camas and Esendal (2006) mentioned that the heritability for plant height, number of branches, seed yield, 1000-seed weight and oil content versus environmental conditions were varied.

ACKNOWLEDGEMENTS

The author acknowledges the Head of the Department of Agronomy, Faculty of Agriculture, Cairo University for providing facilities in the Department. We also wish to express our gratitude to Prof. Dr. R. Shabana and Associate Prof. Dr. Mohamed M. M. Atta for necessary help and assistance.

REFERENCES

- A.O.A.C 2000. Official methods of analysis of A.O.A.C. International. 17th edition by Horwitz, W. Suite (ed.) Vol. (2), chapter (41): 66-68.
- Abu-Hagaza NM, Mahrous N, Mohamed SA, Abd El-Hameed MH. 2009. Response of some promising safflower genotypes to nitrogen levels under drip irrigation in Wadi El-Natroon. *Egyptian J. of Plant breeding*, 13: 183-198
- Abu-Hagaza NM. 1990. Variability and inter-relationships between economic characters in safflower. *Proc. 4th conf. Agron., Cairo, II*, pp. 1-12.
- Amoughin RS, Tobeh A, Somarin SJ. 2012a. Study on the effect of different plant density on some morphological traits and yield of safflower under irrigated and rain-fed planting conditions. *International J. Agron. and Plant Production*, 3(8): 284-290.
- Amoughin RS, Tobeh A, Somarin SJ. 2012b. Effect of plant density on phenology and oil yield of safflower herb under irrigated and rainfed planting systems. *J. Medicinal Plants Res.*, 6(12): 2493-2503.
- Çamaş N, Cüneyt Ç, Enver E. 2007. Seed yield, oil content and fatty acids composition of safflower (*Carthamus tinctorius* L.) grown in northern Turkey conditions. *J. Fac. Agric., OMU*, 22 (1):98-104.
- Camas N, Esendal E, 2006. Estimates of broad-sense heritability for seed yield and yield components of safflower (*Carthamus tinctorius* L.). *Hereditas*, 143: 55-7.
- Elfadl E, Reinbrecht C, Frick C, Claupein W. 2009. Optimizing of nitrogen rate and seed density for safflower (*Carthamus tinctorius*, L.) production under low-input farming conditions in temperate climate. *Field Crops Res.*, 114: 2-13.
- Emami T, Naseri R, Falahi H, Kazemi E. 2011. Response of yield, yield component and oil content of safflower (cv. Sina) to planting date and plant spacing on row in rainfed conditions of western Iran. *American-Eurasian J. Agric. And Environ. Sci.*, 10(6): 947-953.
- Freed RSP, Einensmith S, Gutez S, Reicosky D, Smail VW, Wolberg P. 1989. User's guide to MSTAT-C analysis of agronomic research experiments. Michigan Univ. East Lansing, USA.
- Ghanem SAI, Ash-Shormillesy SMI. 2007. Effect of plant distribution and nitrogen fertilization on yield and photosynthates partitioning of safflower. *Bull. Fac. Agric., Cairo Univ.*, 58:1-10.
- Gomez KA, Gomez AA. 1984. *Statistical Procedures for Agricultural Research* 2nd ed. (Eds. John Willey and Sons), New York, USA.
- Hallauer AR, Miranada JB. 1988. *Quantative genetics in maize breeding* 2nd edition Iowa state Univ. Press, Ames, IA., USA
- Mokhtassi-Bidgoli, A., G. A. Akbari, M. J. Mirhadi, A. R. Pazoki, and S. Soufizadeh (2007). Yield components, leaf pigment contents, patterns of seed filling, dry matter, LAI and LAID of some safflower (*Carthamus tinctorius* L.) genotypes in Iran. *Pakistan J. Biol. Sci.*, 10 (9):1406-1413.
- Muñoz-Valenzuela S, Chanda Musa G, Montoya-Coronado L, Rivera-Rojas VM. 2007. Evaluation of safflower genotypes in Northwest México. *Issues In New Crops and New Uses*. (Ed. J. Janick and A. Whipkey), ASHS Press, Alexandria, VA.
- Osman EBA, Ali A. 2006. Response of some safflower genotypes to modern systems of irrigation in sandy calcareous soils. *Proc.1st Field Crops Conf. Egypt*, pp. 363-371.
- Samanci B, Ozkaynak E. 2003. Effect of planting date on seed yield, oil content and fatty acid composition of safflower (*Carthamus tinctorius*) cultivars grown in the Mediterranean region of Turkey. *J. Agron. and crop Sci.*, 189: 359-360.
- Shahri A, Ganjali HR, Fanayi HR. 2013. Effect of drought on quantitative and qualitative yield of safflower (Goldasht cultivar) in different planting densities. *International J. Agric. and Crop Sci.*, 6(19): 1342-1346.
- Sharifi S, Naderidarbaghshahi M, Golparvar A, Nayerain-Jazy AH. 2012. Effect of plant density on the PAR extinction coefficient and yield of safflower cultivars. *Technical J. Engineering and Applied Sci.*, 2(8): 223-227.
- Sharifmoghaddasi MR, Omid AH. 2009. Determination of optimum row-spacing and plant density in Goldasht safflower variety. *Advances in Environ. Biology*, 3(3): 233-238.
- Vaghar MS, Shamsi K, Kobraee S, Behrooz R. 2014. The effect of planting row interval and plant density on the phenological traits of safflower (*Carthamus tinctorius* L.) under dry land condition. *International J. Biosciences* 4(12): 202-208.
- Weiss EA. 2000. *Oilseed Crops*, 2nd ed., Blackwell Science, Oxford, Chapter: 4, pp. 93-129.
- Yau SK. 2009. Seed rate effect on rainfed and irrigated safflower yield in Eastern Mediterranean. *The Open Agric. J.*, 3, 32-36.